



National Library  
of Medicine PubMed

PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Search

PubMed

for

Go

Clear

☐ Limits

Index

History

Clipboard

About Entrez

Display

Abstract

Save

Text

Order

Add to Clipboard

Entrez PubMed

Overview

Help | FAQ

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Old PubMed

Related Resources

Order Documents

Grateful Med

Consumer Health

Clinical Alerts

☐ 1 : *Cancer Res* 1997 Nov 1;57(21):4882-8

Related Articles, Books

## Regulation of collagenase-3 expression in human breast carcinomas is mediated by stromal-epithelial cell interactions.

Uria JA, Stahle-Backdahl M, Seiki M, Fueyo A, Lopez-Otin C

Departamento de Bioquímica y Biología Molecular, Facultad de Medicina, Universidad de Oviedo, Spain.

Collagenase-3 (MMP-13) is a recently identified member of the human matrix metalloproteinase gene family that is expressed in breast carcinomas and in articular cartilage from arthritic patients. Here, we have studied the cellular origin of this enzyme in breast carcinomas by in situ RNA hybridization, and we found that collagenase-3 is expressed by stromal cells immediately adjacent to epithelial tumor cells but not by the tumor cells themselves; nor is it expressed by the normal breast glandular epithelium. Consistent with this observation, coculture experiments using human fibroblasts and MCF-7 breast cancer cells revealed that conditioned medium from breast cancer cells stimulated the fibroblastic expression of collagenase-3 mRNA. In contrast, no stimulatory effect was observed when medium from fibroblast cells was added to breast cancer cells. These results strongly suggest that transcription of collagenase-3 in stromal cells is activated by diffusible factors released from epithelial breast cancer cells. A survey of a series of cytokines and growth factors known for their ability to induce collagenase-3 expression in human fibroblasts identified interleukin-1alpha and interleukin-1beta as potential candidates for inducing the expression of this MMP gene in breast carcinomas. According to these results, collagenase-3 should be included among the molecular factors that are detected during the stromal reaction to invasive breast cancer and that, by concerted action, may be essential for tumor growth and progression.

PMID: 9354453, UI: 98014576

Display

Abstract

Save

Text

Order

Add to Clipboard